# **Digoxigenin (DIG) Polyclonal Goat Antibody**

Polyclonal goat anti-DIG antibody labeled with our superior CF® dyes and HRP.

# **Product Description**

This is a polyclonal goat anti-DIG antibody labeled with our superior CF® dyes and HRP. This antibody recognizes the digoxigenin (DIG) hapten for detection of DIG conjugated proteins and nucleic acids.

- Available in 6 bright and photostable CF® dyes and HRP
- Suitable for western, immunofluorescence, and immunohistology in FFPE tissues

See our full selection of anti-tag and anti-hapten antibody conjugates.

# Goat Anti-DIG Tag

Conjugation	Ex/Em	Size	Catalog No.
CF®405S	404/431 nm	50 uL (100 ug)	20858-50uL
		250 uL (500 ug)	20858-250uL
CF®488A	490/515 nm	50 uL (100 ug)	20859-50uL
		250 uL (500 ug)	20859-250uL
CF®568	562/583 nm	50 uL (100 ug)	20860-50uL
		250 uL (500 ug)	20860-250uL
CF®594	593/614 nm	50 uL (100 ug)	20861-50uL
		250 uL (500 ug)	20861-250uL
CF®640R	642/662 nm	50 uL (100 ug)	20862-50uL
		250 uL (500 ug)	20862-250uL
CF®680R	680/701 nm	50 uL (100 ug)	20863-50uL
		250 uL (500 ug)	20863-250uL
HRP	N/A	100 uL (100 ug)	20864-100uL

#### Product attributes

Call us: 800-304-5357

Clonality	Polyclonal
Antibody type	Tag Antibody
Concentration	1 mg/mL (HRP conjugate), 2 mg/mL
Host species	Goat
Antibody reactivity (target)	DIG
Antibody/conjugate formulation	Liquid: PBS/50% glycerol/2 mg/mL BSA/0.05% azide
Cross adsorption	Not cross-adsorbed
Secondary/tag antibody applications	ELISA, Flow cytometry, IHC, IF (cells or tissue sections), Western blot

Email: btinfo@biotium.com

View our full selection of Secondary Antibodies, or search our catalog using our Antibody Finder. Alternatively, you can view our secondary antibody product listings with catalog

CF® Dyes offer exceptional brightness and photostability. For more information see our CF® Dye technology page.

### Storage and Handling

numbers.

Liquid format: Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Liquid format antibodies contain 50% glycerol and will not freeze at -20°C.

Lyophilized format: Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Reconstitute antibodies in water using the indicated volumes below:

CF® Dye and biotin conjugates: add 0.5 mL dH<sub>2</sub>O

HRP or DNP conjugates: add 1 mL dH<sub>2</sub>O

Add the indicated volume of water directly to the vial containing the lyophilized antibody and mix gently to dissolve. Store reconstituted antibody at -20°C and protect from light. Aliquot to avoid repeated freeze/thaw cycles. Alternatively, an equal volume of glycerol can be mixed with the reconstituted antibody so that it will remain liquid at -20°C.

Optional: A preservative such as 0.05% sodium azide (final concentration) can be added to CF® Dye and biotin conjugates. Do not add sodium azide to HRP conjugates.

Note: Storage of the antibody for more than a day at final working dilution is not recommended.

CF is a registered trademark of Biotium, Inc.

## References

Download a list of CF® dye references.

This datasheet was generated on September 20, 2024 at 04:34:33 AM. Visit product page to check for updated information before use. Product link: <a href="http://18.235.89.61/product/cf-dye-hrp-dig-tag-polyclonal-goat-antibody/">http://18.235.89.61/product/cf-dye-hrp-dig-tag-polyclonal-goat-antibody/</a>